

# Bactrin

in recurrent urinary tract infection



## attacks pathogens from site to source

Bactrim continues to demonstrate high clinical effectiveness in recurrent urinary tract infections. Bactrim reaches effective levels in urine, serum and renal tissue<sup>1</sup>...the trimethoprim component diffuses into vaginal secretions in bactericidal concentrations<sup>1</sup>...and in the fecal flora, Bactrim effectively suppresses Enterobacteriaceae<sup>1,2</sup> with little resulting emergence of resistant organisms.

\*Due to susceptible strains of indicated org.



Only Valium® (diazepam/Roche)  
is indicated in anxiety  
disorders and as  
an adjunct  
in the relief  
of skeletal  
muscle spasm

**Before prescribing, please consult complete product information, a summary of which follows:**

**Indications:** Management of anxiety disorders, or short-term relief of symptoms of anxiety. Anxiety or tension associated with the stress of everyday life usually does not require treatment with an anxiolytic. Symptomatic relief of acute agitation, tremor, delirium tremens and hallucinosis due to acute alcohol withdrawal; adjunctively in skeletal muscle spasm due to reflex spasm to local pathology; spasticity caused by upper motor neuron disorders; athetosis; stiff-man syndrome; convulsive disorders (not for sole therapy). The effectiveness of Valium (diazepam/Roche) in long-term use, that is, more than 4 months, has not been assessed by systematic clinical studies. The physician should periodically reassess the usefulness of the drug for the individual patient.

**Contraindicated:** Known hypersensitivity to the drug. Children under 6 months of age. Acute narrow angle glaucoma; may be used in patients with open angle glaucoma who are receiving appropriate therapy.

**Warnings:** Not of value in psychotic patients. Caution against hazardous occupations requiring complete mental alertness. When used adjunctively in convulsive disorders, possibility of increase in frequency and/or severity of grand mal seizures may require increased dosage of standard anticonvulsant medication; abrupt withdrawal may be associated with temporary increase in frequency and/or severity of seizures. Advise against simultaneous ingestion of alcohol and other CNS depressants. Withdrawal symptoms similar to those with barbiturates and alcohol have been observed with abrupt discontinuation, usually limited to extended use and excessive doses. Infrequently, milder withdrawal symptoms have been reported following abrupt discontinuation of benzodiazepines after continuous use, generally at higher therapeutic levels, for at least several months. After extended therapy, gradually taper dosage. Keep addiction-prone individuals under careful surveillance because of their predisposition to habituation and dependence.

**Usage in Pregnancy:** Use of minor tranquilizers during first trimester should almost always be avoided because of increased risk of congenital malformations as suggested in several studies. Consider possibility of pregnancy when instituting therapy; advise patients to discuss therapy if they intend to or do become pregnant.

**Precautions:** If combined with other psychotropics or anticonvulsants, consider carefully pharmacology of agents employed; drugs such as phenothiazines, narcotics, barbiturates, MAO inhibitors and other antidepressants may potentiate its action. Usual precautions indicated in patients severely depressed, or with latent depression, or with suicidal tendencies. Observe usual precautions in impaired renal or hepatic function. Limit dosage to smallest effective amount in elderly and debilitated to preclude ataxia or oversedation.

The clearance of Valium (diazepam/Roche) and certain other benzodiazepines can be delayed in association with Tagamet (cimetidine) administration. The clinical significance of this is unclear.

**Side Effects:** Drowsiness, confusion, diplopia, hypotension, changes in libido, nausea, fatigue, depression, dysarthria, jaundice, skin rash, ataxia, constipation, headache, incontinence, changes in salivation, slurred speech, tremor, vertigo, urinary retention, blurred vision. Paradoxical reactions such as acute hyperexcited states, anxiety, hallucinations, increased muscle spasticity, insomnia, rage, sleep disturbances, stimulation have been reported; should these occur, discontinue drug. Isolated reports of neutropenia, jaundice; periodic blood counts and liver function tests advisable during long-term therapy.

**Dosage:** Individualize for maximum beneficial effect.

**Adults:** Anxiety disorders, symptoms of anxiety, 2 to 10 mg b.i.d. to q.i.d.; alcoholism, 10 mg t.i.d. or q.i.d. in first 24 hours, then 5 mg t.i.d. or q.i.d. as needed; adjunctively in skeletal muscle spasm, 2 to 10 mg t.i.d. or q.i.d.; adjunctively in convulsive disorders, 2 to 10 mg b.i.d. to q.i.d. **Geriatric or debilitated patients:** 2 to 2½ mg, 1 or 2 times daily initially, increasing as needed and tolerated. (See Precautions.) **Children:** 1 to 2½ mg t.i.d. or q.i.d. initially, increasing as needed and tolerated (not for use under 6 months).


**Supplied:** Valium® (diazepam/Roche) Tablets, 2 mg, 5 mg and 10 mg—bottles of 100 and 500; Tel-E-Dose® packages of 100, available in trays of 4 reverse-numbered boxes of 25, and in boxes containing 10 strips of 10; Prescription Paks of 50, available in trays of 10.



Roche Laboratories  
Division of Hoffmann-La Roche Inc.  
Nutley, New Jersey 07110

Only Valium® (diazepam/Roche)  
has these two distinct effects

# mind & muscle



-Skeletal  
muscle  
relaxant

-Antianxiety

2-mg, 5-mg, 10-mg  
• scored tablets  
**Valium®**  
diazepam/Roche

Indicated in anxiety disorders  
and as an adjunct in the relief  
of skeletal muscle spasm.

Please see summary  
of product information  
on preceding page.

